

AD-A239 489



(2)

AN IMPROVED FIRST BOREL-CANTELLI LEMMA

BY

DONALD R. HOOVER

DTIC
ELECTE
AUG 14 1991
S D D

TECHNICAL REPORT NO. 446


JULY 31, 1991

PREPARED UNDER CONTRACT
N00014-89-J-1627 (NR-042-267)
FOR THE OFFICE OF NAVAL RESEARCH

Reproduction in Whole or in Part is Permitted
for any purpose of the United States Government

Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA



01 8 13 054

31-07746
|||||

AN IMPROVED FIRST BOREL-CANTELLI LEMMA

BY

DONALD R. HOOVER

TECHNICAL REPORT NO. 446

JULY 31, 1991

Prepared Under Contract

N00014-89-J-1627 (NR-042-267)

For the Office of Naval Research

Herbert Solomon, Project Director

Accession No.	
NTIS	CR-91
DTIC	DA8
Uncontrolled	
Instruction	
By	
D. R. Hoover	
Availability Codes	
Dist	Avail. and/or Special
A-1	

Reproduction in Whole or in Part is Permitted
for any purpose of the United States Government

Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

INTRODUCTION

Let $E_1, E_2, \dots, E_i, \dots$ be an infinite sequence of events. Define $\{E_n \text{ i.o.}\}$ to be the event that an infinite number of the E_i occur. The well known First Borel–Cantelli Lemma states that:

$$\sum_{i=1}^{\infty} P\{E_i\} < \infty \Rightarrow P\{E_n \text{ i.o.}\} = 0 \text{ or equivalently} \quad (1)$$

$$\lim_{m \rightarrow \infty} \sum_{i=m}^{\infty} P\{E_i\} = 0 \Rightarrow P\{E_n \text{ i.o.}\} = 0. \quad (2)$$

In words this lemma states that if the total sum of probability for all events is finite, then the probability of events occurring infinitely often is zero.

The First Borel–Cantelli Lemma gives only a sufficient condition for $P\{E_n \text{ i.o.}\}$ to be zero. A trivial example where the First Borel–Cantelli Lemma conditions are not met yet $P\{E_n \text{ i.o.}\} = 0$ is now given: Let $P\{E_i\} = \frac{1}{i}$ and let $E_i \supset E_{i+1}$ for all i . Since events after E_m occur only if E_m occurs, $P\{E_n \text{ i.o.}\}$ is less than the limit as m goes to infinity of $P\{E_m\}$, which is zero. Yet the summation of probabilities for the individual events is infinite, hence the First Borel–Cantelli condition is not met.

If, however, in this above example, the events E_i were independent then $P\{E_n \text{ i.o.}\}$ would be one. $P\{E_n \text{ i.o.}\}$ changes dramatically even though the probabilities for the individual events are the same. This underscores the importance of event overlap in infinite occurrence. In one case, the overlap was so large that future events did not occur unless previous events had occurred. Since the probability of an event occurring went to zero, so did $P\{E_n \text{ i.o.}\}$. In the other case, the independence of events acted to ensure that future events

would always occur.

In the next section, known overlap between occurrence of adjacent events is incorporated into the First Borel-Cantelli Lemma expanding its application.

IMPROVING THE FIRST BOREL-CANTELLI LEMMA

$$\text{By definition of i.o., } P\{E_n \text{ i.o.}\} = \lim_{m \rightarrow \infty} P\left\{\bigcup_{i=m}^{\infty} E_i\right\}. \quad (3)$$

Or equivalently for any positive integer p ,

$$P\{E_n \text{ i.o.}\} = \lim_{m \rightarrow \infty} \left[P\left\{\bigcup_{i=m}^{m+p-1} E_i\right\} + \sum_{i=m+p}^{\infty} P\left\{E_i \cap \left[\bigcup_{j=m}^{i-1} E_j\right]^c\right\} \right]. \quad (4)$$

Applying (4) with $p = 1$ to the previous example with $E_i \supset E_{i+1}$ gives

$$P\{E_n \text{ i.o.}\} = \lim_{m \rightarrow \infty} \left[P\{E_m\} + \sum_{i=m+p}^{\infty} 0 \right] = \lim_{m \rightarrow \infty} \frac{1}{m} = 0.$$

Similarly applying (3) or (4) with $p = 1$ to the previous example with the E_i being independent gives $P\{E_n \text{ i.o.}\} = \lim_{m \rightarrow \infty} P\left\{\bigcup_{i=m}^{\infty} E_i\right\} = 1$.

The usefulness of (3) and (4) in practice are limited since the right-hand side of (3) and the second term in the right-hand side of (4) involve probabilities of infinite unions. These probabilities may not be calculable.

It often will be possible to calculate probabilities involving up to p elements. When this

is the case, the second term in the right-hand side of (4) can be bounded by a summation of terms involving p events. It sometimes can be shown that this term vanishes as $m \rightarrow \infty$ resulting in the Improved First Borel-Cantelli Lemma (IFBCL).

THE IFBCL:

If for any positive integer p , $\sum_{i=p+1}^{\infty} P\{E_i \cap [\bigcup_{j=1}^{p-1} E_{i-j}]\}^c$ is finite, (5)

it follows that $\lim_{m \rightarrow \infty} P\{\bigcup_{i=m}^{m+p-1} E_i\}$ exists and equals $P\{E_{\infty} \text{ i.o.}\}$. (5.1)

Proof:

From (4), $P\{E_{\infty} \text{ i.o.}\} = \lim_{m \rightarrow \infty} \left[P\{\bigcup_{i=m}^{m+p-1} E_i\} + \sum_{i=m+p}^{\infty} P\{E_i \cap [\bigcup_{j=m}^{i-1} E_j]\}^c \right]$.

But $0 \leq \lim_{m \rightarrow \infty} \left[\sum_{i=m+p}^{\infty} P\{E_i \cap [\bigcup_{j=m}^{i-1} E_j]\}^c \right] \leq \lim_{m \rightarrow \infty} \left[\sum_{i=m+p}^{\infty} P\{E_i \cap [\bigcup_{j=1}^{p-1} E_{i-j}]\}^c \right]$. (6)

The assumptions stated in (5) imply that the limit of the right-most term of (6) is zero, thus the limit of the middle term in (6) is also zero. Hence, this same term vanishes from the right-hand side of (4) as $m \rightarrow \infty$ giving (5.1).

Notes:

(a) For $p = 1$, if one uses standard convention and considers $\left[\bigcup_{j=1}^{p-1} E_{i-j}\right]$ to be ϕ , then

the IFBCL reduces to the Standard First Borel-Cantelli Lemma.

(b) For $p = 1$, under the assumptions in (5), the limit of (5.1) must always be zero. For $p > 1$, the limit of (5.1) can be any number c in $[0,1]$, as is shown by the trivial example where E_i occurs if and only if E_1 occurs for all $i \geq 1$ and the probability E_1 occurs is c .

(c) For $p_2 > p_1$, $\sum_{i=p_1+1}^{\infty} P\{E_i \cap [\bigcup_{j=1}^{p_1-1} E_{i-j}]\} < \infty$ implies that $\sum_{i=p_2+1}^{\infty} P\{E_i \cap [\bigcup_{j=1}^{p_2-1} E_{i-j}]\} < \infty$. Thus, increasing the value of p results in a loosening of the restrictions.

(d) The IFBCL could be generalized by making the condition in (5) that $\sum_{i=2}^{\infty} P\{E_i \cap [\bigcup_{j=1}^{q_i} B_{i,j}]\}$ is finite, where q_i is any positive integer and $B_{i,1}, B_{i,2}, \dots, B_{i,q_i}$ are any events taken from E_1, \dots, E_{i-1} . The result would then be that:

$$P\{E_n \text{ i.o.}\} = \lim_{m \rightarrow \infty} P\left\{\bigcup_{i=1}^{q_m} B_{m,i}\right\}.$$

An Example with $p = 2$

Consider the probability space (F, Ω, P) where Ω is Angular measure in radians on a circle $[0, 2\pi)$, F is Borel Sets and P is Lebesgue Measure/ 2π . Let c be a constant with $0 \leq c \leq 1$. Let

$$r_i = \left[\left[2\pi \cdot \sum_{j=1}^{i-1} \left[\frac{1}{j} - \frac{1}{j+1} + \left(\frac{1}{j+1}\right)^\alpha \right] \right] \right] \text{ MOD } 2\pi \quad \text{for } \alpha > 1,$$

$f_i(w) = I_{\{w \in [r_i, (r_i + 2\pi/i) \text{ MOD } 2\pi] \cup [0, 2\pi c]\}}$ and E_i occur if and only if $f_i(w)$ is 1. Then E_i

is a sequence with $P\{E_i\} \geq \min(1/i, c)$ and $P\{E_i \cap E_{i-1}^c\} \leq (1/i)^\alpha$ for $i \geq 2$. Since $\sum_{i=1}^{\infty} P\{E_i\}$ diverges, the Standard First Borel–Cantelli Lemma may not be used. But $\sum_{i=2}^{\infty} P\{E_i \cap E_{i-1}^c\} \leq \sum_{i=2}^{\infty} (\frac{1}{i})^\alpha$ which, for $\alpha > 1$, is finite. So by the IFBCL with $p = 2$, $P\{E_n \text{ i.o.}\} = \lim_{m \rightarrow \infty} P\{E_m \cup E_{m+1}\} = \lim_{m \rightarrow \infty} (c + \epsilon)$ where ϵ is $O(\frac{1}{m})$ and thus the limit is c .

In this example, the Improved First Borel–Cantelli Lemma compensates for overlap of successive events to give an exact probability for infinite occurrence in a sequence of events with individual probabilities which are too large to enable application of the standard First Borel–Cantelli Lemma.

RELATION TO IMPROVED BONFERRONI INEQUALITIES

The standard First Borel–Cantelli Lemma and improved versions presented here are asymptotic analogs of the first order Bonferroni upper bound and improved forms of it that have recently appeared in the literature. The standard Bonferroni upper bound written for an infinite number of ordered events starting at event m is:

$$\sum_{i=m}^{\infty} P\{E_i\} \geq P\left\{\bigcup_{i=m}^{\infty} E_i\right\}. \quad (8)$$

If the summation on the left-hand side of (8) is finite, then taking the limit as $m \rightarrow \infty$ on both sides of (8) results in (2). Thus the standard First Borel–Cantelli Lemma arises from taking a limit on the standard Bonferroni inequality applied to an infinite number of events with a finite sum of individual probabilities.

The standard Bonferroni upper bound has been improved to incorporate dependency structure for up to p events (Hunter (1976) for $p = 2$, Hoover (1990) for $p > 2$). This bound applied to an infinite number of ordered events starting at event m gives:

$$P\left\{\bigcup_{i=m}^{m+p-1} E_i\right\} + \sum_{i=m+p}^{\infty} P\left\{E_i \cap \left[\bigcup_{j=1}^{p-1} E_{i-j}\right]^c\right\} \geq P\left\{\bigcup_{i=m}^{\infty} E_i\right\}. \quad (9)$$

Taking the limit of the right-hand side of (9) as $m \rightarrow \infty$ gives $P\{E_n \text{ i.o.}\}$. If the second term on the left-hand side of (9) is finite for any m , then it vanishes as $m \rightarrow \infty$ and thus $\liminf_{m \rightarrow \infty} P\left\{\bigcup_{i=m}^{m+p-1} E_i\right\}$ is an upper bound for $P\{E_n \text{ i.o.}\}$. Clearly $\limsup_{m \rightarrow \infty} P\left\{\bigcup_{i=m}^{m+p-1} E_i\right\}$ is a lower bound for $P\{E_n \text{ i.o.}\}$. Thus both limits must equal each other and $P\{E_n \text{ i.o.}\}$. But this is restating the IFBCL using the improved Bonferroni upper bound applied to an infinite number of events.

While Bonferroni's (1936) other inequalities can be extended to produce Borel-Cantelli type lemmas, doing so will usually produce trivial results, such as $P\{E_n \text{ i.o.}\} \geq 0$ or $P\{E_n \text{ i.o.} \leq d\}$ for d some positive integer larger than 1.

REFERENCES

- Bonferroni, C. (1936), *Teoria Statistica Classica Calcolo Delle Probabilita*, *Pubbl. R. 1st. Sugges. Sci. Econ. Comm. Firenze*, 8, 1-62.
- Hoover, D. (1990), Subset Complement Addition Upper Bounds, *Journal of Stat. Plan. and Inf.* 24, 195-202.
- Hunter, D. (1976), An Upper Bound for the Probability of a Union, *J. Appl. Prob.* 13, 597-603.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 446	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) An Improved First Borel-Cantelli Lemma		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL REPORT
7. AUTHOR(s) Donald R. Hoover		6. PERFORMING ORG. REPORT NUMBER
8. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, CA 94305		6. CONTRACT OR GRANT NUMBER(s) N00014-89-J-1627
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics & Probability Program Code 1111		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-042-267
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE July 31, 1991
		13. NUMBER OF PAGES 8
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) First Borel-Cantelli Lemma, Dependency Structure, Improved Bonferroni Bounds		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An improved First Borel-Cantelli Lemma incorporating dependency structure between subsets of events is presented and applied to an example where the First Borel-Cantelli Lemma can't be used. Analogies to improved Bonferroni upper bounds are given.		